



## Prevalence of glucose-6-phosphosphate dehydrogenase deficiency in the newborns in Sanandaj, Iran

**Nahid Ghotbi<sup>1</sup>, Nadia Adnani<sup>2</sup>**

1 Associate Professor, Kurdistan Research Center for Social Determinants of Health, Kurdistan University of Medical Sciences, Sanandaj, Iran

2 Resident, Department of Pediatrics, School of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Iran

### Short Communication

#### Abstract

**BACKGROUND:** Glucose-6-phosphate dehydrogenase (G6PD) deficiency is a genetic disorder and assessment of newborns with or without this deficiency is an essential component in public health evaluation in different countries. Hence, this study was aimed to assess the prevalence of G6PD deficiency in the newborn population in Sanandaj, Iran.

**METHODS:** This is a cross-sectional study on 2016 newborns in Besat Hospital in Sanandaj, Iran, in the year 2006. Three drops of blood were collected from the infants' heels using sterile needles. Then fluorescent spot test was utilized to study the activity of G6PD enzyme.

**RESULTS:** The results of the present study conducted on 2016 neonates showed that 48.80% (984) of them were males and 51.20% (1032) were females. Prevalence of G6PD deficiency in boys and girls were 7.62% and 2.52%, respectively with a male to female ratio of 3:1.

**CONCLUSION:** G6PD deficiency is a gender related condition with a higher frequency among boys' population.

**KEYWORDS:** Glucose-6-Phosphate Dehydrogenase Deficiency, Fluorescent Spot Test, Sex Related

**Date of submission:** 26 Dec 2013, **Date of acceptance:** 09 Feb 2014

**Citation:** Ghotbi N, Adnani N. **Prevalence of glucose-6-phosphosphate dehydrogenase deficiency in the newborns in Sanandaj, Iran.** Chron Dis J 2014; 2(2): 98-101.

### Introduction

Glucose-6-phosphate dehydrogenase (G6PD) is a universally common hereditary disorder. According to World Health Organization (WHO) reports, 2.9% of the world population and 10-15% of the Iranians have G6PD deficiency.<sup>1,2</sup> This disease is more prevalent in Africa, Asia, the Mediterranean, and the Middle East, and approximately 200-400 million people suffer from it all over the world.<sup>3,4</sup>

G6PD enzyme exists in all cells and has a crucial role in providing cell protection during oxidative stress.<sup>5</sup> In individuals with G6PD

deficiency, lifetime of these cells is lower than normal due to oxidation of red blood cell membrane; therefore, hemolysis occurs. Moreover, excessive red cell hemolysis and increased catabolism raise blood bilirubin and jaundice.<sup>6</sup>

Main symptoms of defect in G6PD enzyme include acute hemolytic anemia and classic appearance of favism, neonatal jaundice, and non-spherocytic hemolytic anemia.<sup>7</sup> Neonatal jaundice is one of the most important and remarkable symptoms of this deficiency.<sup>8</sup> An important problem of hyperbilirubinemia is irreversible neurological complications and profound mental retardation in the newborns, which are highly prevalent in Greek, Nigeria, Saudi Arabia, and Southern Iran.<sup>4,9</sup> Individuals

#### Corresponding Author:

Nahid Ghotbi

Email: [ghotbinahid@yahoo.com](mailto:ghotbinahid@yahoo.com)

with this deficiency cannot be blood donors. Moreover, prolonged contact with mild cell hemolytic individuals can cause chronic anemia.<sup>10</sup>

Identifying neonates with this enzyme deficiency is highly significant. Different studies from around the world have considered screening of neonates regarding G6PD enzyme as a very effective factor in decreasing Kernicterus. Recent studies on health education in ethnic Kurds revealed a high prevalence of this medical and health risk factors.<sup>11-13</sup>

However, few studies have been conducted on G6PD in ethnic Kurds as one of the Iranian ethnicities and prevalence of the disease in Sanandaj, Iran, has not been identified yet; therefore, detection of the disease would help reduction of side-effects by consuming food and medication containing antioxidants. Hence, the present study was aimed at identifying prevalence of G6PD dehydrogenase deficiency in neonates in Sanandaj.

### Materials and Methods

This is a cross-sectional study conducted on 2016 neonates born in Besat Hospital in Sanandaj from 2006 to 2007. Permissions were obtained from neonates' parents. Then three drops of blood collected from the infants' heels using sterile needles after recording of the infants' characteristics.

Fluorescent spot test was utilized to study the activity of G6PD enzyme. For this purpose, 100 µl of reagent put in a small container and 10 µl of whole blood containing ethylenediaminetetraacetic acid were taken from the neonates' feet which was added and mixed afterwards. Subsequently, the solution was kept at the room temperature for 20 min, a drop was taken from it using a 20-µl sampler and put on the filter paper. Then the paper was left out to dry. The resulted spot was put under fluorescent light with the wavelength of 365 µm and its reflected light studied, then the collected data were registered. Samples with G6PD enzyme

had the ability to catalyze the chemical reaction, reflect the fluorescent light, and turned into green; however, those with G6PD enzyme deficiency turned into black and could not reflect the light. Data were analyzed in SPSS for Windows (version 16.0, SPSS Inc., Chicago, IL, USA) using chi-square and odds ratio (OR).

### Results

The present study conducted on 2016 neonates including 984 (48.80%) males, and 1032 (51.20%) females. Prevalence of G6PD dehydrogenase deficiency was 5.00%, 7.62% in male infants and 2.52% in female. The ratio of male to female in this particular enzyme was 3.02, in addition, chi-square test proved a statistically significant relationship between G6PD dehydrogenase deficiency and neonates' sex ( $P = 0.0001$ ). The OR was 3.19 (confidence interval 1.98-5.17) (Table 1).

**Table 1. Relationship between sex and glucose-6-phosphate dehydrogenase deficiency, in the male and female newborns of Sanandaj, Iran**

Sex	With G6PD deficiency, n (%)	Without G6PD deficiency, n (%)
Male	75 (74.3)	909 (47.5)
Female	26 (25.7)	1006 (52.5)
Total	101 (100)	1915 (100)

G6PD: Glucose-6-phosphate dehydrogenase

### Discussion

G6PD dehydrogenase deficiency is a sex-related disorder and has different prevalence rates in different countries. In the present study, prevalence of G6PD dehydrogenase deficiency was 5.0%. Its level of prevalence varies in different regions of the country. It has been reported to be 19.3% in Southeastern Iran,<sup>14</sup> 10.9% in Mazandaran,<sup>15</sup> 7.5% in Isfahan,<sup>16,17</sup> 3.2% in Tehran,<sup>18</sup> and 2.1% in Zanjan.<sup>19</sup> In a similar study conducted in Rafsanjan by Alidalaki, G6PD prevalence was 5.0%.<sup>10</sup> Prevalence of G6PD dehydrogenase deficiency was 25.0% in Oman,<sup>20</sup> and 22.0% in Nigeria;<sup>21</sup> despite a lower rate of prevalence of 5.0% and 3.0% in Canada and UK respectively.<sup>22,23</sup> Therefore, G6PD

prevalence in Sanandaj compared to other countries in the world is in the middle. In addition, due to differences in level of prevalence in different parts of Iran from one side and different countries on the other side, it could be concluded that factors such as race, geographical conditions, and weather are responsible for the differences in the number of affected cases.

Prevalence of G6PD dehydrogenase deficiency in boys and girls were 7.62% and 2.52%, respectively with a male to female ratio of 3:1. In Kazemi et al. study, female population outnumbered the disease cases.<sup>18</sup> In a study conducted by Khalesy et al., boys suffered from G6PD dehydrogenase deficiency 5 times more than girls.<sup>24</sup> In Alidalaki et al. study, there was no significant difference between boys and girls regarding the incidence of the disease.<sup>10</sup> In a study conducted in Mazandaran by Ahmadi and Ghazizadeh, it was reported that the incidence of the disease in boys were 3 times more than girls.<sup>25</sup>

Due to sex-related nature of G6PD enzyme deficiency, in different parts of Iran, especially where the disease incidence among girls and boys is the same, differences in levels of prevalence in the two sexes is likely to be related to regional and racial factors. However, it is necessary to conduct related studies further on a larger scale targeting all over Iran. Therefore, it is recommended to screen, especially cord blood of male infants right after birth. It is also recommended that further national studies on the prevalence of G6PD enzyme deficiency be conducted.

### Conclusion

Prevalence of G6PD dehydrogenase deficiency was 5.00%, 7.62% in male infants and 2.52% in female in Sanandaj. Therefore, G6PD dehydrogenase deficiency is a gender related condition with a higher frequency among boys' population.

### Conflict of Interests

Authors have no conflict of interests.

### Acknowledgments

This article was adopted from dissertation of Nadia Adnani. Here, the researchers would like to appreciate all those who provided us with their sincere help and attention particularly personnel of delivery ward and operation room of Besat Hospital, Sanandaj, Iran.

### References

1. Kaplan M, Hammerman C. Glucose-6-phosphate dehydrogenase deficiency and severe neonatal hyperbilirubinemia: a complexity of interactions between genes and environment. *Semin Fetal Neonatal Med* 2010; 15(3): 148-56.
2. Kaplan M, Hammerman C. Glucose-6-phosphate dehydrogenase deficiency: a potential source of severe neonatal hyperbilirubinaemia and kernicterus. *Semin Neonatol* 2002; 7(2): 121-8.
3. Nkhoma ET, Poole C, Vannappagari V, Hall SA, Beutler E. The global prevalence of glucose-6-phosphate dehydrogenase deficiency: a systematic review and meta-analysis. *Blood Cells Mol Dis* 2009; 42(3): 267-78.
4. Segal GB. Enzymatic defects. In: Kliegman RM, Behrman RE, Jenson HB, Editors. *Nelson Textbook of Pediatrics*. 18<sup>th</sup> ed. Philadelphia, PA: Elsevier Science Health Science Division; 2007. p. 2039-42.
5. Abolghasemi H, Mehrani H, Amid A. An update on the prevalence of glucose-6-phosphate dehydrogenase deficiency and neonatal jaundice in Tehran neonates. *Clin Biochem* 2004; 37(3): 241-4.
6. Glucose-6-phosphate dehydrogenase deficiency. WHO Working Group. *Bull World Health Organ* 1989; 67(6): 601-11.
7. Luzzatto L. Glucose 6-Phosphate Dehydrogenase Deficiency. In: Nathan DG, Oski FA, Editors. *Nathan and Oski's Hematology of Infancy and Childhood*. 6<sup>th</sup> ed. Philadelphia, PA: Saunders; 2003. p. 721-42.
8. Usanga EA, Ameen R. Glucose-6-phosphate dehydrogenase deficiency in Kuwait, Syria, Egypt, Iran, Jordan and Lebanon. *Hum Hered* 2000; 50(3): 158-61.
9. Haj Ebrahim Tehrani F, Faghih Zadeh S, Borna H. Necessity of glucose-6-phosphate dehydrogenase activity determination in hyper bilirubinaemia newborns. *Daneshvar Med* 2004; 11(51): 29-32. [In Persian].
10. Alidalaki S, Negahban T, Halakooei M, Sayadi AR, Ahmadi Kohnali J, Esmaeilzadeh JM, et al. Investigation of Glucose-6-phosphate Dehydrogenase Deficiency in Rafsanjan, Autumn 2004. *J Rafsanjan Univ Med Sci* 2007; 6(4): 290-8. [In Persian].

11. Shams Alizadeh N, Mohsenpour B, Ghaderi E, Razaee F, Delavari A. Quality of life in 15-64 years old people in Kurdistan province, western part of Iran. *J Fundam Ment Health* 2010; 12(1): 448-56. [In Persian].
12. Arasteh M, Alizadeh NS, Ghaderi E, Farhadifar F, Nabati R, Gharibi F. Survey of the Prevalence of Sexual Dysfunctions in Kurdish Women. *J Sex Marital Ther* 2013.
13. Yazdanpanah K, Moghimi N, Yousefinejad V, Ghaderi E, Azizi A, Nazem SF. Dyspepsia prevalence in general population aged over 20 in the west part of Iran. *J Pak Med Assoc* 2012; 62(7): 672-6.
14. Pishva N, Amoozgar H. Hyperbilirubinemia Following exchange transfusion with G6PD deficient donor blood. *Iran J Med Sci* 2001; 26: 143-5. [In Persian].
15. Hashemi SA, Zahed Pasha Y, Haji Ahmadi M, Alaei B. Prevalence of G6PD deficiency among primary school students in Amol. *J Babol Univ Med Sci* 2005; 7(1): 52-6. [In Persian].
16. Iranpour R, Akbar MR, Haghshenas I. Glucose-6-phosphate dehydrogenase deficiency in neonates. *Indian J Pediatr* 2003; 70(11): 855-7.
17. Iranpour R, Hashemipour M, Talaei SM, Soroshnia M, Amini A. Newborn screening for glucose-6-phosphate dehydrogenase deficiency in Isfahan, Iran: a quantitative assay. *J Med Screen* 2008; 15(2): 62-4.
18. Kazemi A, Norouzi H, Tohidi Moghaddam M, Naderloo A. Prevalence of Glucose-6-phosphate dehydrogenase (G6PD) deficiency of newborns in Rasol Akram and Ali Asghar hospitals of Tehran. *Iran South Med J* 2013; 16(1): 61-8. [In Persian].
19. Koosha A, Rafizadeh B. Evaluation of neonatal indirect hyperbilirubinaemia at Zanjan Province of Iran in 2001-2003: prevalence of glucose-6-phosphate dehydrogenase deficiency. *Singapore Med J* 2007; 48(5): 424-8.
20. Nair PA, Al Khusaiby SM. Kernicterus and G6PD deficiency-a case series from Oman. *J Trop Pediatr* 2003; 49(2): 74-7.
21. Ogunlesi TA, Dedeke IO, Adekanmbi AF, Fetuga MB, Ogunfowora OB. The incidence and outcome of bilirubin encephalopathy in Nigeria: a bi-centre study. *Niger J Med* 2007; 16(4): 354-9.
22. Sgro M, Campbell D, Shah V. Incidence and causes of severe neonatal hyperbilirubinemia in Canada. *CMAJ* 2006; 175(6): 587-90.
23. Manning D, Todd P, Maxwell M, Jane PM. Prospective surveillance study of severe hyperbilirubinaemia in the newborn in the UK and Ireland. *Arch Dis Child Fetal Neonatal Ed* 2007; 92(5): F342-F346.
24. Khalesy N, Khosravi N, Haghighi M. Prevalence of Glucose-6-phosphate dehydrogenase deficiency in neonates born in Tehran-Iran (2008-09). *J Gorgan Uni Med Sci* 2012; 14(1): 100-5. [In Persian].
25. Ahmadi AH, Ghazizadeh Z. Evaluation of glucose-6-phosphate dehydrogenase deficiency without hemolysis in icteric newborns at Mazandaran province, Iran. *Pak J Biol Sci* 2008; 11(10): 1394-7.